

40. 死亡者死因年齡別 (全國) 大正十一年

Table with 14 columns: 年, 腸窒扶斯, 麻疹, 百日咳, 實及布格魯布, 流行性感胃, 赤痢, 肺結核, 核性他疾, 癌, 脚氣, 腦膜炎, 腦軟化血及腦, 心的疾ノ器質, 急性氣管支. Rows include age groups (0-4, 5-9, etc.) and a total row.

Table with 14 columns: 年, 慢性氣管支, 肺炎及氣管, 胃ノ疾患, 下痢及腸炎, 腎ヲ臟ノ疾ト及氏ノ病, 婦人疾ノ生殖, 產褥熱, 爾及産他ノ疾, 畸形弱及質先天, 老衰, 自殺, 爾ニ依ル外死, 其患ノ他ノ疾, 合計. Rows include age groups and a total row.

41. 死亡者男女及死因別 大正十一年

Table with 16 columns: 地方, 腸窒扶斯, 麻疹, 百日咳, 實布格魯布, 流行性感胃, 赤痢, 肺結核, 男女. Rows list various regions (北海道, 東北區, 關東區, etc.) and a total row.

41. 死亡者男女

及死因別 大正十一年 續

Table with columns for region (地方), cancer (癌), pneumonia (脚氣), meningitis (腦膜炎), cerebral hemorrhage (腦出血及化), heart disease (心臟ノ器質的疾患), acute asthma (急性氣管支炎), and chronic asthma (慢性氣管支炎). Rows list regions from 1 (Hokkaido) to 47 (Tokushima) and a total (總計).

Table with columns for causes of death: pneumonia and bronchitis (肺炎及氣管支肺炎), stomach disease (胃ノ疾患), intestinal disease (下痢及腸炎), kidney and liver disease (腎臟炎及アライト氏病), congenital weakness (畸形及先質), old age (老衰), suicide (自殺), and region number (地方番號). Rows list the same regions as the first table and a total (總計).